

# KINDIFENAC TABLETS

## COMPOTION

### Each E/C tablets

Diclofenac sodium USP 100, 50, 25 mg

### Indications

pain and inflammation in rheumatic disease(including juvenile idiopathic arthritis) and other musculoskeletal disorders;  
mild to moderate pain including dysmenorrhoea; postoperative analgesia;  
migraine; dental pain; fever with discomfort and pain in children;  
post-immunisation pyrexia

### Cautions and contra-indications

- Hypersensitivity against diclofenac
- History of allergic reaction following the use of all NSAIDs
- Third-trimester pregnancy
- Active stomach and/or duodenal ulceration or gastrointestinal bleeding
- Sever insufficiency of the heart
- Sever liver insufficiency
- Sever renal insufficiency
- Hepatic prophyria
- Breast-feeding should be used with caution during breast-feeding

### Side-effects

Gastro-intestinal disturbances including discomfort, nausea, diarrhoea, and occasionally bleeding and ulceration occur  
not to use more than one oral NSAID at a time, and to remember that all NSAIDs non-selective NSAIDs in patients with a history of peptic ulceration.  
Other side-effects include hypersensitivity reactions(particularly rashes, angioedema, and bronchospasm, headache, dizziness, nervousness, depression, drowsiness, insomnia, vertigo, hearing disturbances such as tinnitus, photosensitivity, and haematuria. Blood disorders have also occurred. Fluid retention may occur (rarely precipitating congestive heart failure); blood pressure may be raised.

### Asthma

Any degree of worsening of asthma may be related to the ingestion of NSAIDs

### Dose

Max. total daily dose by any route 150mg

- By mouth, 75–150 mg daily in 2–3 divided dose
- Child 1-2 years, juvenile arthritis, by mouth 1-3mg/kg daily in divided doses (25mg e/c tablets)

**The tablet should be swallowed without chewing, with some liquid during or immediately after meals**

**Presentation** : Pack of 30 tablets in 3 PVC-ALU sheets with leaflet.

**Storage** : store bellow 30°C& Protected from light.

---

This is medicament  
-Medicament is product affects your health and its consumption contrary  
To instructions is dangerous for you.  
-Follow strictly the Doctors prescription, the method of use the instructions of the pharmacist who sold the medicament .  
-The doctor and the pharmacist are experts in medicines ,their benefits and risks .  
-Don't by yourself interrupt the period of treatment prescribed .  
-Don't repeat the same prescription without consulting your doctor .  
KEEP ALL MEDICAMENTS OUT OF REACH OF CHILDREN

---

AL-KINDI CO. For Pharmaceutical Ind. / Iraq



شركة الكندي للصناعات الدوائية- العراق